

Mepilex® Border Post-Op

Healing uninterrupted

Mepilex Border Post-Op is designed to meet the clinical challenges and criteria set up for post-operative dressings by leading surgeons in the field¹. No patient deserves less than an advanced wound dressing.



Stays on
for up to
7 days¹⁻³

Supports early patient mobility²⁻⁴

By being conformable, flexible and minimising risk of painful skin damage.

Helps reduce the risk of SSIs²⁻⁴

By minimising skin damage and number of potentially contaminating dressing changes.

Reduces dressing related costs^{2,3}

By minimising number of dressing changes.



Avance® Solo

Consistent pressure. For constant care.

Avance Solo is a portable single-patient-use negative pressure system². It is indicated for use on closed surgical incisions, where it aims at supporting undisturbed healing and improved patient mobility.

Mepilex Border Post-Op meets all expert requirements²⁻⁴

- ✓ Stays on well
- ✓ Flexible
- ✓ Showerproof¹
- ✓ No dead space
- ✓ Protection of the skin
- ✓ Good absorption




-125mmHg⁵

A continuous -125mmHg negative pressure is the clinical standard and proven effective in reducing incidence of dehiscence, seroma, haematoma.



CFM Technology™⁶⁻⁹

A combination of controlled air flow, multilayer dressing and distal canister, enabling the Avance® Solo NPWT system to deliver continuous regulated negative pressure.



Safetac® silicone adhesive^{2,3,10-13}

Proven to minimise trauma to wound site, and pain to the patient during dressing removal.

Identify the patient specific risk profile¹⁴



Risk Assessment Tool by Mölnlycke®

High risk

High risk patients are recommended Negative Pressure Wound Therapy for closed surgical incisions (ciNPT), like Avance Solo.

Elevated risk

Elevated risk patients are recommended advanced wound dressings. ciNPT should be considered if cost benefits allows.

Low to moderate risk

Low to moderate risk patients are recommended advanced wound dressings for closed incisions, like Mepilex Border Post-Op.

Learn more about the Risk Assessment Tool and download an editable pdf for using the tool in practice to help make evidence-based decisions in your clinic.

Add
QR-code

References:

1. Sandy-Hodgetts, K., et al. Incision care and dressing selection in surgical wounds: Findings from a series of international meetings. *Wounds International*, 2022.
2. Bredow J. et al. Evaluation of Absorbent Versus Conventional Wound Dressing. A Randomized Controlled Study in Orthopedic Surgery. *Deutsche Arzteblatt International*, 2018.
3. Zarghooni K. et al. Is the use of modern versus conventional wound dressings warranted after primary knee and hip arthroplasty? *Acta Orthopaedica Belgica*, 2015.
4. Beele H. et al. A prospective randomized controlled clinical investigation comparing two post-operative wound dressings used after elective hip and knee replacement; Mepilex® Border Post-Op versus Aquacel® Surgical. *International Journal of Orthopaedic and Trauma Nursing*, 2020.
5. Data on file (15).
6. Data on file(10).
7. Data on file (18).
8. Data on file (23).
9. Data on file (17).
10. Silverstein P. et al. An open, parallel, randomized, comparative, multicenter study to evaluate the cost-effectiveness, performance, tolerance, and safety of a silver-containing soft silicone foam. *Journal of Burn Care and Research*, 2011.
11. Gee Kee E.L. et al. Randomized controlled trial of three burns dressings for partial thickness burns in children. *Burns*, 2014.
12. David F. et al. A randomised, controlled, non-inferiority trial comparing the performance of a soft silicone-coated wound contact layer [Mepitel One] with a lipidocolloid wound contact layer [UrgoTul] in the treatment of acute wounds. *International Wound Journal*, 2017.
13. Patton M.L. et al. An open, prospective, randomized pilot investigation evaluating pain with the use of a soft silicone wound contact layer vs. a traditional dressing and staples on split thickness skin grafts as a primary dressing. *Journal of burn care & research*, 2013.
14. SSERA Group. Surgical patient population risk assessment: The simplified SSERA assessment model. *Wounds International*, 2023.



Find out more at www.molnlycke.com

Mölnlycke Health Care AB, Box 13080, Gamlestadsvägen 3C, SE-402 52 Göteborg, Sweden. Phone +46 31 722 30 00. The Mölnlycke, Mepilex and Avance trademarks, names and logotypes are registered globally to one or more of the Mölnlycke Health Care group of companies.

© 2023 Mölnlycke Health Care AB. All rights reserved. HQIM004544



Dedicated to new beginnings

Every patient deserves the best possible start after surgery. Mölnlycke® Incision Care Solutions help you deliver high-quality outcomes that bring patients back to home and health.